

CH302 Worksheet 0: A Little Thermo Review


1. What two processes (a.k.a. path functions) can transfer internal energy between a system and its surroundings? What symbols are used for these variables?
2. Write a good definition for a state function.
3. Name some state functions.
4. The first law states that the value of which state function is conserved in an isolated system? What **two** symbols are used for this state function?
5. What is an isolated system? Name the most obvious example of an isolated system (hint: big).
6. What **inequality** is often associated with the second law of thermodynamics? What does it mean in plain English?
7. What **equality** is often associated with the second law of thermodynamics? What does it mean in plain English?
8. What does the third law of thermodynamics state?
9. How many translational, rotational and vibrational modes, respectively, does C<sub>2</sub>H<sub>2</sub> have?
10. What would be the total internal energy associated with the vibrational motion of 1 molecule of C<sub>2</sub>H<sub>4</sub>? What about 1 mole of C<sub>2</sub>H<sub>4</sub>?
11. In the list of elements below, mark (circle, underline, etc.) all of the elements that are not shown in their standard state.  
Cdiamond(s)Ca(s)B<sub>2</sub>(s)Na(s)Fe(s)Hg(s)  
Br<sub>2</sub>(l)Mo(s)H(g)He(g)Xe(g)Rb<sub>2</sub>(s)  
Cd(l)As(s)N<sub>2</sub>(l) O<sub>2</sub>(l)Si<sub>60</sub>(s)F<sub>2</sub>(g)
12. Write the standard formation reactions for the following chemical species  
NH<sub>3</sub>(g)  
  
Fe<sub>2</sub>O<sub>3</sub>(s)  
  
O<sub>2</sub>(l)

O<sub>3</sub>(g)


NH<sub>2</sub>OH(s)

13. Assume we want to use a bomb calorimeter to determine the specific heat capacity of an unknown liquid. We use 3 L of the unknown liquid and perform a known reaction that releases 400 kJ of heat. We measure an initial and final temperature of 25 °C and 28.7 °C, respectively. If the heat capacity of the calorimeter is 85 J·K<sup>-1</sup>, and the density of the liquid is 2.34 g·mL<sup>-1</sup>, what is the specific heat capacity of the unknown liquid?

14. Given the following data:



15. Consider the reaction below and approximate the value of the work function at room temperature. (Note: this doesn't require a calculator.)



16. If a system at -272 °C absorbs 545 J of heat, what is its change in entropy?

17. Ammonia (the common name for NH<sub>3</sub>, the odor of windex and cat urine), has a ΔH<sup>°</sup><sub>vap</sub> = 23.35 kJ·mol<sup>-1</sup> and a ΔS<sup>°</sup><sub>vap</sub> = 97.43 J·mol<sup>-1</sup>·K<sup>-1</sup>. What is the normal boiling point of ammonia expressed in centigrade?

18. Consider the reaction below:

HNO<sub>3</sub>(l) + H<sub>2</sub>(g) ↔ H<sub>2</sub>O(l) + NO<sub>2</sub>(g)  
Using the provided table values, calculate ΔG<sup>°</sup><sub>rxn</sub> if it is performed under standard conditions.

	ΔH <sup>°</sup> <sub>f</sub> (kJ·mol <sup>-1</sup> )	ΔS <sup>°</sup> <sub>m</sub> (J·mol <sup>-1</sup> ·K <sup>-1</sup> )
HNO <sub>3</sub> (l)	-174.1	156
H <sub>2</sub> (g)	not provided	131
H <sub>2</sub> O(l)	-285.8	70
NO <sub>2</sub> (g)	33.2	240

19. Rank the following compounds in terms of decreasing standard molar entropy:  $\text{CO(s)}$ ,  $\text{CO}_2(\text{g})$ ,  $\text{CO}_2(\text{l})$ ,  $\text{CO}(\text{l})$ .

20. Assuming you have one mole of each of the following, rank them in terms of decreasing value of the term  $W$  in the Boltzmann equation:  $\text{HF}$ ,  $\text{NaF}$ ,  $\text{ClF}$ ,  $\text{F}_2$ .