Spring 2006 CH302 Worksheet 1b: Examples of Simple Thermodynamics Multiple Choice Problems

- 1. Which of the following is a correct statement concerning the Second Law of Thermodynamics?
 - 1. The free energy of a system is temperature dependent.
 - 2. Entropy of a system increases in the phase change from a liquid to a gas.
 - 3. Energy cannot be created nor destroyed.
 - 4. The entropy in the universe is conserved.

2. 75 g of a potato chips are burned in a calorimeter that contains 2 liters of water

initially at 297K. After the combustion, the temperature rises 12° C. How much heat is evolved per gram of potato chip burned? The heat capacity of the calorimeter is 200 J/°C? The density of water is 1.0 g/ml. The specific heat of water is 4.18 J/g°C.

1. 7.1 kJ/g 2. 100.3 kJ/g 3. 102.7 kJ/g 4. 1.37 kJ/g

- 3. What is the change of enthalpy associated with the combustion of one mole of ethylene?
 - 1. 0 kJ 2. -1323 kJ 3. +1323 kJ 4. -3230 kJ 5. +3230 kJ

4. For the combustion reaction of acetaldehyde (C2H4O)

$$2C_2H_4O + 5O_2 \rightarrow 4CO_2 + 4H_2C$$

assume all reactants and products are gases and calculate the $\Delta H^{0}rxn$ using the following bond energy values:

C-C BE = 611 kJ/mol C-H BE = 413 kJ/mol O=O BE = 498 kJ/mol C=O BE = 799 kJ/mol H-O BE = 463 kJ/mol 1. -1080 kJ/mol

1. -1080 kJ/mol 2. +1080 kJ/mol 3. 0 kJ/mol 4. -2303 kJ/mol 5. +2303 kJ/mol

$$C_{2}H_{4} + 3O_{2} \rightarrow 2CO_{2} + 2H_{2}O$$

5. For the reaction

$$3H_2(g) + N_2(g) \rightarrow 2NH_3(g)$$

find the approximate value for the work done at 300 K.

1. -5.0 kJ 2. -2:5 kJ 3. 2.5 kJ 4. 5.0 kJ

6. Heat flow is considered negative when heat flows (into, out of) a system; work is considered positive when work is done (by, on) a system.

1. out of; by
2. into; by
3. out of; on
4. into; on

7. Which of the following processes results in an increase in the system entropy?

- 1. cleaning up from the party while your parents are out of town
- 2. getting dressed in the morning
- 3. making ice cubes
- 4. pouring salt on an icy bridge
- 5. memorizing the eight question types on the first quiz
- 8. For the exothermic combustion of a hydrogen balloon:

$$2H_2 + O_2 --> 2 H_2O_{(g)}$$

what can you say about the spontaneity?

- 1. Always spontaneous because ΔS in formation H₂O_(g) is negative.
- 2. Always spontaneous because ΔS in formation $H_2O_{(g)}$ is positive.
- 3. Spontaneous at higher temperature because ΔS in formation $H_2O_{(g)}$ is negative
- 4. Spontaneous at lower temperature because ΔS in formation $H_2O_{(g)}$ is negative.
- 5. Spontaneous at higher temperature because ΔS in formation $H_2O_{(g)}$ is positive.
- 6. Spontaneous at lower temperature because ΔS in formation $H_2O_{(g)}$ is positive.